

On Lange's Conjecture

Montserrat Teixidor i Bigas,

Mathematics Department, Tufts University, Medford MA 02155, U.S.A., mteixido@tufts.edu

temporary address DPMMS, 16 Mills Lane, Cambridge CB2 1SB, England. teixidor@dpmms.cam.ac.uk

Let C be a generic curve of genus g defined over an algebraically closed field of characteristic zero. Denote by $U(n, d)$ the moduli space of stable vector bundles of rank n and degree d on C . Let E be an element of $U(n, d)$. Consider the set of subbundles E' of E of rank n' . Define the integer $s_{n'}(E) = n'd - n \max\{\deg E'\}$. Then, E is semistable if and only if $s_{n'}(E) \geq 0$ for all n' . On the other hand, a generic $E \in U(n, d)$ must satisfy $s_{n'}(E) \geq n'(n - n')(g - 1)$ (cf [L] Satz p.448), while for any E , $s_{n'}(E) \leq n'(n - n')g$ (cf [L] Satz p.448, [M-S]).

One can then stratify $U(n, d)$ according to the value of $s_{n'}$: define

$$U_{s,n'}(n, d) = \{E \in U(n, d) | s_{n'}(E) \leq s\}, 0 \leq s \leq n'(n - n')g$$

These loci are closed in $U(n, d)$ because the function $s_{n'}(E)$ is upper semicontinuous on E . It is not clear though whether the loci are non-empty. The most natural approach towards the study of $U_{s,n'}(n, d)$ is to consider families of extensions of vector bundles E'' by vector bundles E' . Here E'' is assumed to have rank $n - n'$ and degree $d - d'$. If $\mu(E') < \mu(E'')$, one expects the generic such extension to be stable. One could even restrict attention to the subloci $\bar{U}_{s,n'}(n, d)$ of vector bundles that can be written as an extension with both E', E'' semistable. A moduli count shows that the dimension of the set of such extensions is $d_{s,n'} = n^2(g - 1) + 1 + s - n'(n - n')(g - 1)$. Hence, one expects $\dim U_{(s,n')}(n, d) = d_{s,n'}$ if $0 < s \leq n'(n - n')(g - 1)$ and $U_{(s,n')}(n, d) = U(n, d)$ if $s \geq n'(n - n')(g - 1)$. Lange (cf [L]) conjectured that $\bar{U}_{s,n'}(n, d)$ should be non-empty for $0 < s \leq n'(n - n')(g - 1)$. The conjecture had been proved so far for vector bundles of rank two (cf [L, N]) and for vector bundles of any rank if $0 < s < \min(n', n - n')(g - 1)$ and in a few very special cases (cf [B, B, R]). The method used in these papers consists in showing that the set of unstable extensions fill a space of dimension smaller than $d_{(s,n')}$. After this work was completed, we received a preprint from H. Lange and L. Bambrila-Paz proving results similar to ours.

We take a different approach which continues our study in [T3]. We work with a curve generic in the sense of moduli. We construct explicitly stable vector bundles of degree d and rank n that have subbundles of rank n' and degree d' in the suitable range for a special type of reducible curve. We then show that this is enough in order to prove the result for the generic curve. Our method also allows us to clarify what happens in the so far grey area $n'(n - n')(g - 1) \leq s \leq n'(n - n')g$. The situation is as follows: subbundles of rank n' and degree d' exist on the generic E as soon as $s \geq n'(n - n')(g - 1)$, $s \equiv n'd(n)$. Our result is

(0,1) Theorem. *Let C be a generic curve of genus $g \geq 2$ defined over a field of characteristic zero. Fix integers $n, d, 0 < n' < n, s \equiv n'd(n)$. Define $d' = (n'd - s)/n$.*

i) *If $0 \leq s \leq n'(n - n')(g - 1)$, then $\bar{U}_{s,n'}(n, d)$ is non-empty and there are elements $E \in \bar{U}_{s,n'}(n, d)$ that have only a finite number of subbundles of rank n' and maximal degree.*

ii) *Assume $n'(n - n')(g - 1) \leq s$. Let E be a generic vector bundle on C and Q a generic point on C . Then E can be written as an extension $0 \rightarrow E' \rightarrow E \rightarrow E'' \rightarrow 0$, $\text{rk } E' = n', \deg E' = d'$ and both E', E'' semistable. The set $A = A(n, d, E)$ of subbundles of E of rank n' and degree d' is non-empty of dimension $s - n'(n - n')(g - 1)$ and the map*

$$\begin{array}{ccc} \pi_Q : & A & \rightarrow \mathbf{G}(n', E_Q) \\ & E' & \rightarrow E'_Q \end{array}$$

is either onto or has generic finite fibers.

Note: for elliptic curves, the result is essentially correct (see (2,8)) except that stability must be replaced by semistability. This is to be expected as a "generic" vector bundle on an elliptic curve is only semistable

and not stable if rank and degree are not coprime (see [Tu] for an alternative approach to moduli spaces of vector bundles on elliptic curves).

The paper is structured as follows: in section 1 we gather a few technical results. These will allow us to reduce the proof of the Theorem for the generic curve to the proof in the case of a special reducible curve. Specifically, we are going to deal with the curve that can be obtained by gluing at a point a generic curve of genus one less with an elliptic curve. Section 2 proves (a strenghtening of) the Theorem for the case of an elliptic curve. In addition to providing the first step of the induction, it will give us the working tool to allow us to treat the tail of our reducible curve. This section makes heavy use of Atiyah's classical paper [A]. We need here characteristic zero (see the proof of (2,8)) although the results are likely to hold in any characteristic. The final section will prove the Theorem for the reducible curve under the assumption that it holds for the generic curve of smaller genus.

Acknowledgements : This work has been carried out during a visit to the Pure Mathematics Department at the University of Cambridge. The author is a member of the Europroj group "Vector Bundles on Curves". The interest in this question was spurred by a talk given by Barbara Russo at the Europroj meeting 96. Thanks are also due to Peter Newstead for pointing out an incorrect definition of $s_{n'}(E)$.

1 Reduction to the case of a reducible curve.

We mainly use standard notation. We gather below a few definitions, including some that have already been introduced above.

(1,1) Notation. As above $U(n, d)$ denotes the moduli space of stable vector bundles on a curve. When the curve is reducible, stability is understood with respect to a polarisation (a_i) (cf.[S]). $U_{s,n'}(n, d)$ denotes the set of stable vector bundles that have a subbundle of rank n' and degree d' with $n'd - nd' = s$. When there is some danger of confusion, we shall add reference to the curve as $U_{s,n'}(n, d)(C)$.

$\bar{U}_{s,n'}(n, d)$ denotes the set of stable vector bundles that have a stable subbundle of rank n' and degree d' with $n'd - nd' = s$ and the quotient is also stable.

$A(n', d', E)$ will denote the set of subbundles of rank n' and degree d' of a vector bundle E on the curve C . When C is reducible, we shall often replace the degree by a multidegree (or degree of the restriction to the different components). If the curve is C_k, C^k instead of C , we shall write the sub or superindex k in A as well.

$\mathbf{G}(n', V)$ denotes the Grassman variety of linear subspaces of dimension n' of a vector space V of dimension n .

$\pi_P : A(n', d', E) \rightarrow \mathbf{G}(n', E_P)$ denotes the (rational) map that sends a subbundle to its fiber at a (generic) point $P \in C$

(1,1) Lemma. *Let C be a projective connected reduced curve having only nodes as singularities. Let $0 \rightarrow E' \rightarrow E \rightarrow E'' \rightarrow 0$ be an exact sequence of vector bundles. Assume that the restriction to each component of C of each of the vector bundles above is semistable and that $\mu(E') < \mu(E)$. Then $h^0(E''^* \otimes E') = 0$.*

Proof: A non-zero section of $E''^* \otimes E'$ is equivalent to a non-zero map $f : E'' \rightarrow E'$. As the restriction of E to each component C_i is semistable, $\mu(E'_{|C_i}) \leq \mu(E''_{|C_i})$. Moreover, as $\mu(E') < \mu(E)$, there is at least one component C_{i_0} such that $\mu(E'_{|C_{i_0}}) < \mu(E''_{|C_{i_0}})$. By the semistability of the bundles involved, $f_{|C_{i_0}} = 0$. As C is connected, given any component C_{i_1} , there exists a chain $C_1, \dots, C_t$ such that $C_1 = C_{i_0}, C_t = C_{i_1}$ and C_i intersects $C_{i-1}, i = 2 \dots t$. We prove by induction on t that $f_{|C_t} \equiv 0$. If $t = 1$, we know this already. Assume $f_{|C_{t-1}} = 0$. Then the map $f_{|C_t}$ vanishes at a point P in the intersection of C_{t-1} and C_t . Hence it gives rise to a map $\bar{f}_{|C_t} : E''(P)_{|C_t} \rightarrow E'_{|C_t}$. As $\mu(E''(P)_{|C_t}) > \mu(E'_{|C_t})$ the semistability of the bundles implies that $\bar{f}_{|C_t} = 0$. Hence $f_{|C_t} = 0$. This concludes the induction and hence the proof of the Lemma.

The following proposition and section 3 deal with vector bundles on reducible nodal curves. There is a notion of stability for torsion-free sheaves on such a curve introduced by Seshadri. We refer the reader to [S] Ch.VII-VIII, where he constructs a moduli spaces for such objects. Here we recall how to determine a vector bundle on such a curve.

Let C be a nodal curve with irreducible components C_i and nodes P_j . The restriction of a vector bundle E to each C_i is a vector bundle E_i . If P_j is in the intersection of C_{i_1} and C_{i_2} , there is a "gluing"

isomorphism of the vector spaces fiber at P_j : $E_{i_1, P_j} \cong E_{i_2, P_j}$. Conversely, a set of such data determines a vector bundle. A torsion-free sheaf is determined by giving isomorphisms of suitable quotients of the fibers $E_{i_1, P_j}/V_{i_1, j} \cong E_{i_2, P_j}/V_{i_2, j}$.

(1,3) Proposition. *Let C^0 be a tree-like curve (or curve of compact type). Let*

$$0 \rightarrow E'^0 \rightarrow E^0 \rightarrow E''^0 \rightarrow 0$$

be an exact sequence of vector bundles on C^0 . Assume that the restriction of each vector bundle to each component of C^0 is semistable and stable on at least one of them. Assume also that the restriction of the vector bundles to any destabilising rational component is trivial (i.e. of the form $\mathcal{O}^n$). Denote by n, d, n', d' the rank and degree of E^0, E'^0 respectively, $s = n'd - nd'$.

i) Assume $0 < n'd - nd' \leq n'(n - n')(g - 1)$, and E^0 has only a finite number of subbundles of rank n' and degree d' . Then, $\bar{U}_{s, n'}(n, d)(C)$ is non-empty for the generic curve C of genus g and the generic element has only a finite number of subbundles of rank n' and degree d' .

ii) Assume $n'(n - n')(g - 1) \leq s$ and $\dim A^0(n', d', E^0) = n'd - nd' - n'(n - n')(g - 1)$. Then the generic element E in $U(n, d)(C)$ has a semistable subbundle of rank n' and degree d' with semistable quotient and $\dim A(n', d', E) = n'd - nd' - n'(n - n')(g - 1)$.

Proof: This proof is inspired in Lange's construction of $\bar{U}_{s, n'}$ (cf. [L] section 4) adapted to the case of a family and is essentially contained in [T3]. We include it here for the sake of completeness.

Claim: The set of vector bundles that are trivial on rational components and satisfy the stability conditions stated above can be identified to an open set of the moduli space of torsion-free sheaves on the stable model $\tilde{C}$ of C^0 : Consider a vector bundle on C^0 , that is trivial on a rational component, There is a canonical isomorphism between the fibers at the two nodes P_1, P_2 on the rational component induced by the pair of isomorphisms $H^0(\mathbf{P}^1, \mathcal{O}^n) \rightarrow \mathcal{O}_{P_i}^n$ $i = 1, 2$. This induces a natural gluing isomorphism for the vector bundle on the curve with this component blown down. Proceed in this way until all components are blown down. Conversely, given a vector bundle (say $\tilde{E}$) on $\tilde{C}$, one can build a vector bundle on C^0 by taking trivial bundles $\mathcal{O}^n$ on the additional rational components. The gluings at the nodes are canonically determined by those of $\tilde{E}$.

Conversely, we can replace C^0 by a curve C'^0 with chains of rational components added between some of the nodes of C^0 (i.e. a semistable curve with same stable model as C^0 a little bit more unstable than C^0).

Consider a family $\mathcal{C} \rightarrow S$ parametrised say by the spectrum of a discrete valuation ring whose central fiber is C^0 and whose generic fiber is a non-singular curve C of genus g . From the reasoning above, adding rational components to C^0 does not alter the hypothesis. Hence, we can make base changes and normalise if needed.

We now need the existence of moduli spaces parametrising vector bundles of the type E'^0, E^0, E''^0 on $\mathcal{C}$. In fact we need suitable coverings of these moduli spaces with Poincare families. Denote by $(C_i)_{i \in I}$ the irreducible components of C^0 . Choose a polarisation $(a_i)_{i \in I}$ for C^0 . This is the choice of rational numbers ("weights") $a_i > 0, \sum a_i = 1$. Consider moduli spaces of torsion-free sheaves on $\mathcal{C} \rightarrow S$ of ranks $n', n, n - n'$ and degrees $d', d, d - d'$ such that the restriction to the generic fiber is semistable and the restriction to the special fiber is (a_i) -semistable. These can be shown to exist as in Seshadri's proof for a single curve by replacing points by sections of the family. Alternatively, one can use the existence of a moduli space for torsion-free sheaves over the moduli space $\bar{\mathcal{M}}_g$ of stable curves constructed by Pandharipande([P]). Our curves are only semistable and not necessarily stable but our vector bundles are trivial on rational components. By blowing down these rational components, we can identify the open set of these bundles with an open set of the moduli space of vector bundles on the stable model of the curve as explained above. Take then in the standard way suitable coverings over which Poincare bundles exist. Denote by $\mathcal{U}', \mathcal{U}, \mathcal{U}''$ these moduli spaces, by $\mathcal{M}', \mathcal{M}, \mathcal{M}''$ their coverings and by $\mathcal{E}', \mathcal{E}, \mathcal{E}''$ the corresponding Poincare bundles. The given vector bundles E'^0, E^0, E''^0 are not necessarily (a_i) -stable and therefore may not be represented in these families. In order to deal with this situation, we need to recall some results about stability of vector bundles on tree-like curves (see [T1], [T2]).

Assume a vector bundle on C^0 is (a_i) -semistable. One can choose an ordering of the components of C^0 in such a way that the complement of the components chosen beforehand is connected. Then the degree on

each component lies on intervals of length n in the real line determined by d and the degree in the previous components. We shall call such a multidegree an (a_i) -allowable multidegree. The converse is not true in general: a vector bundle with an (a_i) -allowable multidegree is not necessarily stable. However if E is a vector bundle with an (a_i) -allowable multidegree, the restriction to each component is semistable and at least one of them is stable, then E is stable. One can choose a multidegree on C^0 such that for every divisor D^0 on C^0 with this multidegree, $E^0(D^0)$ has its multidegree in these intervals. Everything remains valid replacing E^0 by say E'^0, E''^0 or replacing C^0 by $\tilde{C}$.

As C^0 is of compact type, one can choose a divisor $\mathcal{D}$ on $\mathcal{C}$ with support on C^0 whose restriction to C^0 has a given preassigned degree. By the normality of $\mathcal{C}$, $\mathcal{D}$ corresponds to a line bundle $\mathcal{L}$. Similarly, there are line bundles $\mathcal{L}', \mathcal{L}''$ such that $E'^0 \otimes \mathcal{L}'|_{C^0}, E''^0 \otimes \mathcal{L}''|_{C^0}$ are (a_i) -semistable. Then $\mathcal{M}', \mathcal{M}, \mathcal{M}''$ with Poincaré bundles $\mathcal{E}' \otimes \mathcal{L}'^{-1}, \mathcal{E} \otimes \mathcal{L}^{-1}, \mathcal{E}'' \otimes (\mathcal{L}'')^{-1}$ parametrise families of vector bundles containing E'^0, E^0, E''^0 . Moreover, there are open sets of $\mathcal{M}', \mathcal{M}, \mathcal{M}''$ such that the restriction of the corresponding vector bundle to each component of C^0 is semistable. Denote by q, p', p'' the projections of $\mathcal{C} \times_S \mathcal{M}' \times_S \mathcal{M}''$ on the three factors. Consider

$$\mathcal{F} = R^1(p' \times p'')_*[(q \times p'')^*(\mathcal{E}'' \otimes (\mathcal{L}'')^{-1})^* \otimes (q \times p')^*(\mathcal{E}' \otimes (\mathcal{L}')^{-1})].$$

From (1,2) and Riemann-Roch's Theorem, $\mathcal{F}$ is a vector bundle of rank $n'(n - n')(g - 1) + n'd - nd'$ over $\mathcal{M}' \times \mathcal{M}''$. Then, $\mathbf{P}(\mathcal{F})$ parametrises extensions of vector bundles and there is a universal family (cf [L2])

$$0 \rightarrow (id \times \pi)^*(\mathcal{E}' \otimes \mathcal{L}'^{-1}) \rightarrow \tilde{\mathcal{E}} \rightarrow (id \times \pi)^*(\mathcal{E}'' \otimes \mathcal{L}''^{-1}) \rightarrow 0$$

where π denote pull-backs to $\mathbf{P}(\mathcal{F})$. Using the (a_i) -allowability of the multidegrees of the endterms of this sequence, one can check the (a_i) -allowability of the central term. As (semi)-stability is an open condition and the restriction of E^0 to each component of C^0 is semistable, the same is true for the generic extension. Hence the generic extension is (a_i) -stable. There is then a rational map $\mathbf{P}(\mathcal{F}) \rightarrow U$. If $s \leq n'(n - n')(g - 1)$, our assumption implies that the fibers of the map are finite. Hence the image has dimension $n'^2(g - 1) + 1 + (n - n')^2(g - 1) + 1 + n'(n - n')(g - 1) + n'd - nd' - 1 = n^2(g - 1) + 1 + s - n'(n - n')(g - 1)$ as claimed. If $s \leq n'(n - n')(g - 1)$, our assumption implies that the fibers of the map have dimension at most $s - n'(n - n')(g - 1)$. Hence, the map is generically onto as claimed.

Remark. The condition that the restriction of the vector bundles to one of the components of C^0 is stable could be weakened. All we need in fact is that the vector bundle one obtains on the whole curve is stable by some polarisation (up to tensoring each restriction with a line bundle).

2 Subbundles of vector bundles on elliptic curves.

In this section C will always be an elliptic curve. A vector bundle is said to be indecomposable if it is not the direct sum of smaller rank subbundles. Up to the choice of a point P in C , there is a canonically determined indecomposable vector bundle $E(n, d)$ of rank n and degree d on C (cf. [A]). When $d = 0$, $E(n, 0)$ is the only indecomposable vector bundle that has sections. Every indecomposable vector bundle on C is of the form $E(n, d) \otimes L$ for some line bundle L of degree zero. Hence $Pic^0(C) \cong C$ can be taken as a parameter space for the family of indecomposable vector bundles on C . Denote by $\mathcal{L}$ the Poincare bundle of degree zero on $C \times C \cong C \times Pic^0(C)$. Then $\mathcal{E}(n, d) = p_2^*(E(n, d)) \otimes \mathcal{L}$ is a Poincare bundle on $C \times C$ (where p_2 denotes projection onto the second factor).

Every vector bundle on C is a direct sum of indecomposable vector bundles. If we fix a decomposition type, say

$$(2, 1)d^j, n^j > 0, j = 1 \dots k, \sum_j d^j = d, \sum_j n^j = n,$$

one can take as parameter space for vector bundles with this type of decomposition the k^{th} product of the curve C . The corresponding Poincare bundle is

$$(2, 2)\mathcal{E}((n^j, d^j)) = \oplus [p_{k+1}^*(E((n^j, d^j))) \otimes p_{j, k+1}^*(\mathcal{L})].$$

Here again p denote projections and the subindices denote the factors we are projecting onto.

(2,3) Lemma. *Let E, E' be indecomposable vector bundles on C of ranks n, n' and degrees d, d' respectively. Then, $E'^* \otimes E$ is a direct sum of indecomposable vector bundles all of them with the same slope.*

Proof: From [A] p.434 Corollary iv), $E(n', d')^* = E(n', -d')$. If $(n, d) = (n', d') = 1$, the result is then Lemma 33 of [A]. If $(n, d) = h, (n', d') = h', n = \bar{n}h, n' = \bar{n}'h'$ then by [A] Lemma 24, $E(n, d) = E(h, 0) \otimes E(n_1, d_1)$, $E(n', d') = E(h', 0) \otimes E(n'_1, d'_1)$. From [A] Lemma 21, $E(h, 0) \otimes E(h', 0)$ is a direct sum of indecomposable vector bundles of degree zero. Then, the result follows using [A] Lemma 33 applied to $E(\bar{n}, \bar{d}) \otimes E(\bar{n}', \bar{d}')$ and [A] Lemma 24 again.

(2,4) Lemma. *Let E be an indecomposable vector bundle of rank n and degree d . Fix integers $n' > 0, d'$. Consider all possible decomposition types for a vector bundle E' of rank n' and degree d' : $d_i^j, n_i^j > 0, j = 1 \dots k_i, \sum_j d_i^j = d', \sum_j n_i^j = n'$. Let X_i be the set of points in $C \times \dots \times C$ representing vector bundles E'_i such that there is a map from E' to E that is non-zero on each indecomposable summand $E_i'^j$ of E'_i . There is only a finite number of X_i . Define $\mathbf{Hom}_i = p_{1 \dots k_i}^*[\mathcal{E}((n_i^j, d_i^j))^* \otimes p_{k_i+1}^* E]_{|X_i}$. Then, $\mathbf{Hom}_i$ is a vector bundle*

$$(2, 5) \dim X_i + rk \mathbf{Hom}_i \leq n'd - nd' + h^0(E_i'^* \otimes E_i)$$

where E'_i is a generic point of X_i . Define $(n', d') = h', n' = \bar{n}'h', d' = \bar{d}'h'$ and $\bar{X}$ as the component corresponding to the decomposition of E' into direct sum of h' subbundles of degree $\bar{d}'$ and rank $\bar{n}'$. Then on $\bar{X}$ (2,5) is an equality except may be in the case when $n'd - nd' = 0, h' > 0$. Equality in (2,5) implies that all the summands of E' have the same slope. If $(n, d) = 1$ or $n'd - nd' > 0$, equality in (2,5) occurs only on $\bar{X}$.

Proof: We can write $E = E(n, d) \otimes L$. A vector bundle of degree d_i^j and rank n_i^j is of the form $E(n_i^j, d_i^j) \otimes L_i^j$ for a degree zero line bundle L_i^j . A generic point E'_i of X_i is of the form $E'_i = \oplus_j (E(n_i^j, d_i^j) \otimes L_i^j)$. From (2,3), $(E(n_i^j, d_i^j) \otimes L_i^j)^* \otimes E(n, d) \otimes L$ is a direct sum of indecomposable vector bundles all of them of slope $(n_i^j d - nd_i^j)/n_i^j n$. If $n_i^j d - nd_i^j < 0$, then $(E(n_i^j, d_i^j) \otimes L_i^j)^* \otimes (E(n, d) \otimes L)$ has no nontrivial sections (use for instance [A] p.434 Corollary iv) and [A] lemma 15). Hence X_i is empty. Therefore, we need only consider a finite number of decomposition types corresponding to $n_i^j d - nd_i^j \geq 0$ for all possible choices of j .

From (2,3) above and [A] Lemma 15, if $n_i^j d - nd_i^j > 0$

$$h^0([E(n_i^j, d_i^j) \otimes L_i^j]^* \otimes [E(n, d) \otimes L]) = n_i^j d - nd_i^j.$$

Assume now $n_i^j d - n d_i^j = 0$. Write

$$(n, d) = h, (n_i^j, d_i^j) = h_i^j, n = h\bar{n}, d = h\bar{d}, n_i^j = h_i^j \bar{n}_i^j, d_i^j = h_i^j \bar{d}_i^j.$$

Then

$$E(n, d) = E(h, 0) \otimes E(\bar{n}, \bar{d}), E(n_i^j, d_i^j) = E(\bar{n}_i^j, \bar{d}_i^j) \otimes E(h_i^j, 0)$$

As $0 = n_i^j d - n d_i^j = h h_i^j (\bar{n}_i^j \bar{d} - \bar{n} \bar{d}_i^j)$ and $(\bar{n}, \bar{d}) = 1$, it follows that $\bar{n} | \bar{n}_i^j$. Similarly, $\bar{n}_i^j | \bar{n}$. Hence, $\bar{n}_i^j = \bar{n}, \bar{d}_i^j = \bar{d}$. Then

$$(E(n_i^j, d_i^j) \otimes L_i^j)^* \otimes (E(n, d) \otimes L) = E(h_i^j, 0) \otimes E(h, 0) \otimes E(\bar{n}, \bar{d}) \otimes E(\bar{n}, \bar{d})^* \otimes L \otimes (L_i^j)^{-1}.$$

Using [A] Lemma 22, $E(\bar{n}, \bar{d}) \otimes E(\bar{n}, \bar{d})^* = \oplus L_k$ where the L_k are the line bundles of order dividing $\bar{n}$. Hence, from [A] Lemma 21,

$$E(n_i^j, d_i^j)^* \otimes E(n, d) = [E(|h - h_i^j| + 1, 0) \oplus E(|h - h_i^j| + 3, 0) \oplus \dots \oplus E(h + h_i^j - 1, 0)] \otimes \oplus (L_k L(L_i^j)^{-1})$$

Hence the space of maps from $E(n_i^j, d_i^j)$ to $E(n, d)$ is non-empty if and only if $L_i^j = L L_k$ for some k . This means that there is only a finite number of choices for L_i^j . For this finite number of choices, the dimension of the space of maps from $E(n_i^j, d_i^j) \otimes L_i^j$ to $E(n, d)$ is $\min(h, h_i^j)$.

Therefore, $h^0(E_i'^* \otimes E)$ is constant for E_i' on a fixed X_i . Hence, $\mathbf{Hom}_i$ is a vector bundle of rank $h^0(E_i'^* \otimes E)$.

Write $\epsilon_i^j = 0$ if $n_i^j d - n d_i^j > 0$, $\epsilon_i^j = 1$ if $n_i^j d - n d_i^j = 0$. Then

$$\begin{aligned} \dim X_i + rk \mathbf{Hom}_i &= \dim X_i + \sum (n_i^j d - n d_i^j) + \sum \epsilon_i^j \min(h, h_i^j) = \\ &= (k_i - \sum \epsilon_i^j) + n' d - n d' + \sum \epsilon_i^j \min(h, h_i^j). \end{aligned}$$

We now compute the dimension of the space of automorphisms of E_i' . From the description of E_i' ,

$$E_i'^* \otimes E_i' = \oplus_j [E(n_i^j, d_i^j)^* \otimes E(n_i^j, d_i^j)] \oplus \oplus_{j_1 \neq j_2} [(E(n_i^{j_1}, d_i^{j_1}) \otimes L_i^{j_1})^* \otimes (E(n_i^{j_2}, d_i^{j_2}) \otimes L_i^{j_2})]$$

Write $(n_i^j, d_i^j) = h_i^j$ and $n_i^j = h_i^j \bar{n}_i^j, d_i^j = h_i^j \bar{d}_i^j$. Then $E(n_i^j, d_i^j) = E(\bar{n}_i^j, \bar{d}_i^j) \otimes E(h_i^j, 0)$ (cf[A], Lemma 24). From [A] Lemma 22, $E(\bar{n}_i^j, \bar{d}_i^j) \otimes E(\bar{n}_i^j, \bar{d}_i^j)^* = \oplus L_i^{j_k}$ where the $L_i^{j_k}$ are the linebundles on C of order dividing $\bar{n}_i^j$. From [A] Lemma 21,

$$E(h_i^j, 0) \otimes E(h_i^j, 0)^* = E(1, 0) \oplus E(3, 0) \oplus \dots \oplus E(2h_i^j - 1, 0).$$

Hence, $h^0(\text{End} E(n_i^j, d_i^j)) = h_i^j$.

If $n_i^{j_2} d_i^{j_1} - n_i^{j_1} d_i^{j_2} > 0$, then, using (2,3),

$$h^0((E(n_i^{j_2}, d_i^{j_2}) \otimes L_i^{j_2})^* \otimes (E(n_i^{j_1}, d_i^{j_1}) \otimes L_i^{j_1})) = n_i^{j_2} d_i^{j_1} - n_i^{j_1} d_i^{j_2}$$

If $n_i^{j_2} d_i^{j_1} - n_i^{j_1} d_i^{j_2} = 0$, then reasoning as above, there is only a finite number of choices of $L_i^{j_2}$ for which the space of maps to the component corresponding to j_1 is non-zero. For a generic point of $\bar{X}$ the L_i^j are generic except when $n' d - n d' = 0$. So, for a generic point of $\bar{X}$, all these spaces are zero except may be in the case when $n' d - n d' = 0$. Hence,

$$h^0(E_i'^* \otimes E_i') \leq \sum_j h_i^j + \sum_{j_1 < j_2} |n_i^{j_2} d_i^{j_1} - n_i^{j_1} d_i^{j_2}|.$$

with equality for the generic E in $\bar{X}$ if $n' d - n d' > 0$. As $h_i^j \geq 1$ for all j , $\dim X_i + rk \mathbf{Hom}_i \leq n' d - n d' + h^0(E_i'^* \otimes E_i')$ as stated. Moreover, if equality occurs, then $n_i^{j_2} d_i^{j_1} - n_i^{j_1} d_i^{j_2} = 0$ for all pairs j_1, j_2 . Also, $h_i^j = 1$

if $\epsilon_i^j = 0$. The first condition implies that all summands of E'_i have the same slope d'/n' . If $d'/n' \neq d/n$, then $h_i^{j_1} = h_i^{j_2} = 1$. This together with the equality above implies $d_i^{j_1} = d_i^{j_2}$, $n_i^{j_1} = n_i^{j_2}$. Hence, we are dealing with a point of $\bar{X}$. Similarly, if $(d, n) = 1$, then $\min(h, h_i^j) = h = 1$ and equality implies that $h_i^j = 1$. We conclude as before that we are dealing with a point of $\bar{A}$.

(2,5) Lemma. *Let $d, n, d', n', 0 < n' < n$ be integers, $h = (n, d), d = \bar{d}h, n = \bar{n}h$. Let $E = \oplus E_i$ be a vector bundle written as the direct sum of h generic indecomposable vector bundles of rank $\bar{n}$ and degree $\bar{d}$. There is a variety $A = A(n', d', E)$ parametrizing the set of subbundles of E of rank n' and degree d' . The variety A is a finite union of components all of them of dimension at most $n'd - nd'$ (and is empty if $n'd - nd' < 0$). The component $\bar{A}$ corresponding to an E' in $\bar{X}$ (as defined in (2,4)) has precisely this dimension.*

Proof: Consider the possible splitting type of a bundle E' of rank n' and degree d' . As an immersion cannot vanish on any direct summand of E' , only a finite number of these decomposition types are possible (cf.(2,4)). With the notations of (2,4), E' corresponds to a point in some X_i . Let X be the (disjoint) union of the components X_i in (2,4) such that the generic map of a point in X_i to E is an immersion. Let $\mathcal{G}$ be the Grassmanian bundle of $\dim n'$ subspaces of the fibers of E . There is then a rational map from $\mathbf{P}(\cup \mathbf{Hom}_i)$ to $\mathcal{G}$. Denote by A its image. Lange proved that A can be identified to the image subvariety in $\mathcal{G}$ (cf.[L], proof of Lemma 4.2). If two maps from the same sheaf E' to E differ in an automorphism of E' , they give rise to the same subbundle of E . Conversely, if two immersions of vector bundles have the same subbundle as image, then, the two subbundles are isomorphic and the immersions differ in an automorphism of the subbundle. Hence, from (2,4), $\dim A \leq n'd - nd'$ with equality only on the component $\bar{A}$ corresponding to $\bar{X}$ if $n'd - nd' > 0$.

When $\bar{n}\bar{d} - \bar{d}'\bar{n} = 0$, then $\bar{n} = \bar{n}', \bar{d} = \bar{d}'$. It is then easy to see that E' must be the direct sum of h' of the summands appearing in the decomposition of E . Hence, the result is obvious.

It remains to show that $\bar{A}$ (and hence A) is non-empty by proving that the generic map from a point of $\bar{X}$ to E is injective. This is clear if $n'd - nd' > 0$.

Assume now $n'd - nd' > 0$ and the generic map from a point E' in $\bar{X}$ to E is not injective. Then the kernel K is a subsheaf of E . As E is torsion-free, K is a vectorbundle of rank $n'' < n'$ and degree d'' . The map $f : E' \rightarrow E$ can be factored through the canonical projection $E' \rightarrow E'/K$ and f is determined by K and the immersion $E'/K \rightarrow E$. As E is torsion-free, so is E'/K and hence E'/K is a vector bundle. Then,

$$\begin{aligned} n'd - nd' &= \dim \bar{A}(n', d', E) \leq \dim A(n'', d'', E') + \dim A(n' - n'', d' - d'', E) \leq \\ &\leq n''d' - n'd'' + (n' - n'')d - n(d' - d''). \end{aligned}$$

Hence,

$$(2, 5, 1)(d - d')/(n - n') \leq d''/n''.$$

On the other hand, by the semistability of E' (see for example [Tu] Appendix A),

$$(2, 5, 2)d''/n'' \leq d'/n'$$

The hypothesis $n'd - nd' > 0$ is equivalent to

$$(2, 5, 3)d'/n' < (d - d')/(n - n').$$

As (2,5,1),(2,5,2),(2,5,3) are incompatible, we obtain a contradiction and the result is proved.

(2,6) Lemma. *With the notations of (2,5), let E be a direct sum of vector bundles all of the same slope. Let E' be a generic point in $\bar{X}$ and $f : E \rightarrow E'$ a generic immersion. Define E'' by the exact sequence*

$$0 \rightarrow E' \rightarrow E \rightarrow E'' \rightarrow 0$$

Then E'' is a direct sum of indecomposable bundles all of the same slope.

Proof: There is a one to one correspondence between immersions of subbundles E' of E of rank n' and degree d' and quotients of E of rank $n - n'$ and degree $d - d'$. Hence, from (2,4), the set of such quotients has dimension $n'd - nd'$.

We show first that E'' is a vector bundle. If this were not the case, the immersion $E' \rightarrow E$ would factor through an immersion $E' \rightarrow \tilde{E}' \rightarrow E$ with $\tilde{E}'/E'$ a torsion sheaf. Write $\tilde{d}' = \deg \tilde{E}' = d' + k, k > 0$. There is then an exact sequence $0 \rightarrow E' \rightarrow \tilde{E}' \rightarrow \mathbf{C}^k \rightarrow 0$, where $\mathbf{C}^k$ is a skyscraper sheaf of length k . The dimension of the set of such extensions is at most $n'k$. The dimension of the set of maps $\tilde{E}' \rightarrow E$ is at most (from (2,5)) $n'd - n\tilde{d}'$. Hence we find a bound $n'd - nd' = \dim \bar{A}(n', d', E) \leq n'k + n'd - n(d' + k) = n'd - nd' - k(n - n')$. As $n \geq n'$, this implies $k = 0$ and so $\tilde{E}' = E'$.

Dualizing the exact sequence defining E'' , we obtain a one to one correspondence between the set of quotients E'' and subbundles of E^* of rank $n - n'$ and degree $-(d - d')$. From (2,4), the dimension of this set is at most $(n - n')(-d) - n(-(d - d')) = nd' - n'd$ and equality implies that E'' is of the stated type. This finishes the proof.

(2,7) Lemma. *Let E be an indecomposable vector bundle with slope $\mu \geq 1$. Then E is generically generated by global sections.*

Proof: We use induction on the rank n . For $n=1$, every line bundle of strictly positive degree on an elliptic curve has sections. So, the result follows. Assume the result for vector bundles of rank at most $n - 1$. Let E be an indecomposable vector bundle of rank n and degree $d \geq n$. Write $d = kn + \epsilon, 0 \leq \epsilon \leq n - 1, k \geq 1$. Let L_k be a line bundle of degree k . If $\epsilon > 0$, $h^0(E \otimes L_k^{-1}) > 0$. If $\epsilon = 0$, there is one L_k such that $h^0(E \otimes L_k^{-1}) > 0$. On the other hand, for any line bundle L of degree at least $k + 1$, $E \otimes L^{-1}$ has negative degree and hence no sections. There is then an exact sequence

$$0 \rightarrow L_k \rightarrow E \rightarrow E'' \rightarrow 0$$

Here E'' has no torsion (otherwise E would have a line subbundle of degree higher than k). From (2,6), E' is a direct sum of vector bundles all of the same slope $(d - k)/(n - 1) = k(n - 1) + \epsilon/(n - 1) \geq k$. By induction assumption, E'' is generically generated by global sections. As $h^0(E) = h^0(L_k) + h^0(E'')$, the result follows.

(2,8) Proposition. *Let $E, A, \bar{A}$ be as in (2,5). Let P, Q be generic points of C . Denote by $\mathbf{G}(n', E_P)$ the Grassman variety of n' -subvector spaces of the fiber of E at P . Consider the rational map*

$$\begin{array}{ccc} \pi_P : & A & \rightarrow \mathbf{G}(n', E_P) \\ & F & \rightarrow F_P \end{array}.$$

Define π_Q similarly.

i) If $0 \leq n'd - nd' < n'(n - n')$ the image by π_P of each component of A has dimension at most $n'd - nd'$ and one component with image of this dimension and finite fibers does exist.

ii) If $n'(n - n') \leq n'd - nd'$, then π_P is surjective and so the fibers of π_P have dimension $n'd - nd' - n'(n - n')$.

iii) If $n'd - nd' \leq 2n'(n - n')$, then the product map $\pi_P \times \pi_Q$ has generic finite fibers.

iv) If $2n'(n - n') \leq n'd - nd'$, then the product map $\pi_P \times \pi_Q$ is onto.

Proof: Assume first $0 \leq n'd - nd' \leq n'(n - n')$. From (2,3) the dimension of every component of A is at most $n'd - nd'$. We only need to show that the restriction of π_P to $\bar{A}$ has finite fibers. We shall prove that the differential of π_P is injective.

The tangent space to A at a point corresponding to an immersed vector bundle E' is the space of infinitesimal deformations of E' inside E . These are parametrised by $H^0(E'^* \otimes E/E')$. The tangent space to a grassmanian of subspaces of a fixed dimension of a space V at a point corresponding to a subspace V' is given by $\text{Hom}(V', V/V')$. In our case, $V' = E'_P, V = E_P$. We need to prove then that the map

$$d\pi_P : H^0(E'^* \otimes (E/E')) \rightarrow E'^* \otimes (E_P/E'_P)$$

is injective when E' is generic in $\bar{A}$ and P is generic in C . By (2,6), E/E' is a direct sum of indecomposable vector bundles all of the same slope. Then, (2,3) implies that $E'^* \otimes E/E'$ is a direct sum of indecomposable

vector bundles all of the same slope. The assumption $0 \leq n'd - nd' < n'(n - n')$ can be written as $0 \leq (d - d')n' - (n - n')d' < n'(n - n')$. Hence the slope of each indecomposable summand of $E'^* \otimes E/E'$ is positive and less than one. From [A] Lemma 15 ii), there is an immersion

$$H^0(E'^* \otimes E/E') \otimes \mathcal{O}_C \rightarrow E'^* \otimes E/E'$$

Taking the fiber at the point P , the map will remain injective. This proves i).

Assume now $n'(n - n') \leq n'd - nd'$. We need to show now that the map $d\pi_P$ above is surjective. From (2,3) and (2,7) $E'^* \otimes E/E'$ is generically generated by global sections. This is equivalent to ii).

If $n'd - nd' < 2n'(n - n')$, from (2,3) $E'^* \otimes E/E'(-P - Q)$ is a direct sum of indecomposable vector bundles of negative slope. Hence it has no sections. Consider the exact sequence

$$0 \rightarrow E'^* \otimes E/E'(-P - Q) \rightarrow E'^* \otimes E/E' \rightarrow (E'^* \otimes E/E')_{P+Q} \rightarrow 0$$

As $H^0(E'^* \otimes E/E'(-P - Q)) = 0$, taking homology we obtain iii).

Under the hypothesis in iv), $h^0(E'^* \otimes E/E'(-P - Q)) = h^0(E'^* \otimes E/E') - n'(n - n')$. Hence, $H^0(E'^* \otimes E/E')$ maps onto $(E'^* \otimes E/E')_{P+Q}$. This proves iv).

3 Proof of the result for the reducible curve.

(3,1)Lemma. *Let $n', n, 0 < n' < n$ be integers. Let V_1, V_2 be vector spaces of dimension n . Let A_1, A_2 be subvarieties of the Grassmannians $\mathbf{G}(n', V_1), \mathbf{G}(n', V_2)$ of dimensions k_1, k_2 respectively. Consider the set X of linear isomorphisms of V_1 to V_2 that identify some element in A_1 with some element in A_2 .*

- i) $\dim X \leq k_1 + k_2 + n^2 - n'(n - n')$
- ii) If A_2 is a finite set, then $\dim X = k_1 + k_2 + n^2 - n'(n - n')$.
- iii) If $\dim X = k_1 + k_2 + n^2 - n'(n - n')$, then for the generic $\varphi \in X$, there is only a finite number of pairs $(V'_1, V'_2) \in A_1 \times A_2$ such that $\varphi(V'_1) = V'_2$.
- iv) If $X = \text{Is}(V_1, V_2)$, for a generic $\varphi_0 \in \text{Is}(V_1, V_2)$, $\dim\{(V'_1, V'_2) \in A_1 \times A_2 \mid \varphi_0(V'_1) = V'_2\} = k_1 + k_2 - n'(n - n')$

Proof: Consider the set of triples

$$\tilde{X} = \{(\varphi, V'_1, V'_2) \in \text{Is}(V_1, V_2) \times A_1 \times A_2 \mid \varphi(V'_1) = V'_2\}$$

Denote by p_0, p_1, p_2 the projections from $\text{Is}(V_1, V_2) \times \mathbf{G}(n', V_1) \times \mathbf{G}(n', V_2)$ to the corresponding factors. The fiber of $p_1 \times p_2$ over any point in $\mathbf{G}(n', V_1) \times \mathbf{G}(n', V_2)$ has dimension $n^2 - n'(n - n')$. Hence, the inverse image $\tilde{X}$ of $A_1 \times A_2$ by $p_1 \times p_2$ has dimension $k_1 + k_2 + n^2 - n'(n - n')$. As $X = p_0(\tilde{X})$, i) follows. Moreover, if there is equality in the bound for the dimension of X , then the fibers of $p_0 : \tilde{X} \rightarrow X$ are finite. Hence, iii) follows. If $A_2 = \{V'_2\}$ consists of a single point, then p_0 induces a bijection $\tilde{X} \rightarrow X$, the inverse being given by $\varphi \rightarrow (\varphi, (\varphi)^{-1}(V'_2), V'_2)$. This proves ii).

The assumption in iv) means that $p_0 : \tilde{X} \rightarrow \text{Is}(V_1, V_2)$ is onto. Hence the generic fiber has dimension $\dim \tilde{X} - \dim \text{Is}(V_1, V_2)$. This is iv).

(3,2) Proposition. *For $g \geq 2$ let C be a curve obtained by gluing a generic point P_g in a curve C_g of genus g satisfying (0,1) with a point P_1 on a generic elliptic curve C_1 . For $g = 1$, glue two generic elliptic curves C_1, C'_1 at generic point P_1, P'_1 . Fix integers $n, d, n' < n, s \equiv n'd(n)$. Define $d' = (n'd - s)/n$. Then, there exists a polarisation (a_i) of C such that*

- i) If $0 < s \leq n'(n - n')(g - 1)$, there exists a vector bundle E on C that can be written as an extension

$$0 \rightarrow E' \rightarrow E \rightarrow E'' \rightarrow 0$$

with $\text{rk} E' = n', \deg E' = d'$, E has only a finite number of subbundles of rank n' and degree d' and no subbundles of rank n' and larger degree. Moreover, the restrictions of E, E', E'' to C_1 are semistable and

the restrictions to C_g are stable if $g > 1$. If $g = 1$, the restrictions to C'_1 are only semistable but the resulting vector bundles on C are stable for some polarisation.

ii) Assume $n'(n - n')(g - 1) \leq s$. Let $E \in U(n, d, (a_i))(C)$ be generic. The set A of subbundles of E of rank n' and degree d' is non-empty of dimension $s - n'(n - n')(g - 1)$ and the map

$$\begin{array}{ccc} \pi_Q : & A & \rightarrow \mathbf{G}(n', E_Q) \\ & E' & \rightarrow E'_Q \end{array}$$

is either onto or has generic finite fibers.

Proof: We assume $g \geq 2$. We shall deal with the case $g = 1$ at the end.

1) Assume that $0 < s \leq n'(n - n')(g - 1)$. Using (0,1), one can choose a vector bundle E_g on C_g of degree d with $s_{n'}(E_g) = s$ and only a finite number of vector bundles of rank n' and maximal degree. Take as vector bundle E_1 on C_1 a direct sum of n line bundles of degree zero. Glue a direct sum of n' of these line bundles with one of the maximal rank n' subbundles of E_g . Then the resulting vector bundle E has a subbundle that gives the desired value s . It cannot have higher degree subbundles because the highest degree for a vector bundle of E_1 is zero and we are assuming $s_{n'}(E_g) = s$. Moreover, the number of subbundles of maximum degree is finite because there are only a finite number of possibilities for the restriction of E' to C_1, C_g . By (0,1), the conditions on stability are satisfied on C_g . By construction, they are satisfied on C_1 .

2) Assume now $n'(n - n')(g - 1) \leq s \leq n'(n - n')g, s \equiv n'd(n)$. Take a direct sum of n generic linebundles of degree zero on C_1 . Take a generic vector bundle E_g on C_g of degree d . Consider the set A_g of subbundles of E_g of rank n' and degree d' . By (0,1), the image $\pi_{g,Q}(A_g)$ in $\mathbf{G}(n', E_{g,Q})$ has dimension $a_g(d') = s - n'n''(g - 1)$. Consider the set A_1 of subbundles of E_1 of rank n' and degree zero. Then A_1 is finite and so is $\pi_1(A_1)$. Consider the set X of gluings of E_{g,P_g} with E_{1,P_1} that identify one direction in $\pi_{g,Q}(A_g)$ with one subspace in $\pi_1(A_1)$. By (3,1), the dimension of this set is $a_g + 0 + n^2 - n'(n - n') = s + n^2 - n'(n - n')g$. Choose a generic element in X to glue E_{g,P_g} to E_{1,P_1} . The resulting E has a subbundle of rank n' and degree d' . By (0,1), the conditions on stability are satisfied on $C_g, g > 1$. By (2,8) and (2,6), they are satisfied on C_1 . By [T1] step 2, they give rise to a stable E .

We need to show that E has only a finite number of subbundles of rank n' and degree d' and that it does not have subbundles of higher degree.

Assume it had a subbundle of higher degree. Denote by $\bar{d}_g$ and $\bar{d}_1$ the degrees of the restriction to C_g and C_1 respectively. Then $\bar{d}_1 \leq 0$. Hence

$$n'd - n\bar{d}_g = n'd - n(\bar{d}_g + \bar{d}_1) + n\bar{d}_1 < s + n\bar{d}_1 \leq s \leq n'(n - n')g.$$

Moreover, as E_g is generic, $n'd - n\bar{d}_g \geq n'(n - n')(g - 1)$ (cf [L] p.448 Satz). Hence, the induction assumption (in part ii)) of the Theorem applies and the image $\pi_{P_g}(A(n', \bar{d}_g, E_g))$ in $\mathbf{G}(n', E_{P_g})$ is $a_g(\bar{d}_g) = n'd - n\bar{d}_g - n'(n - n')(g - 1) < n'(n - n')$.

As $n'd - n\bar{d}_g \geq n'(n - n')(g - 1)$,

$$n\bar{d}_1 = n'd - n\bar{d}_g - (n'd - n\bar{d}_g - n\bar{d}_1) > n'd - n\bar{d}_g - s \geq n'(n - n')(g - 1) - s \geq -n'(n - n')$$

By (2,8) $\pi_{P_1}(A_1(n', \bar{d}_1, \bar{E}_1))$ has dimension $a_1 = -n\bar{d}_1$. Now, $a_g(\bar{d}_g) + a_1(\bar{d}_1) = n'd - n\bar{d}_g - n\bar{d}_1 - n'(n - n')(g - 1) < s - n'(n - n')(g - 1) \leq n'(n - n')$. By (3,1), the dimension of the set of gluings that glue one subspace in $\pi_{g,Q}(A_g(n', \bar{d}_g, E_g))$ with one subspace in $\pi_1(\bar{A}_1(n', \bar{d}_1, E_1))$ has dimension at most

$$a_g(\bar{d}_g) + a_1(\bar{d}_1) + n^2 - n'(n - n')g = \dim X.$$

Hence the generic element in X does not glue any pair of these subspaces and no such subbundle of E exists.

Assume now that $\bar{d}_g, \bar{d}_1$ are chosen in some other form so that $n'd - n\bar{d}_g - n\bar{d}_1 - n'(n - n')(g - 1) = s$. One can repeat the proof above except that the strict inequalities are now equalities. By (0,1), the fibers of $\pi_{g|A_g(n', \bar{d}_g, E_g)}$ are finite. By (2,8), the fibers of $\pi_{1|A_1(n', \bar{d}_1, E_1)}$ are finite. By (3,1), a generic gluing in X glues at most a finite number of subspaces in $\pi_g(A_g(n', \bar{d}_g, E_g))$ with subspaces in $\pi_1(A_1(n', \bar{d}_1, E_1))$. This shows that the number of subbundles obtained in this way is finite.

3) Assume now $n'(n-n')g \leq s$. Choose d_1, d'_1 such that $n'd_1 - nd'_1 = n'(n-n')$. This is possible because the greatest common divisor of n, n' divides also $n'(n-n')$.

Write $h = (n, d_1), n = h\bar{n}, d_1 = h\bar{d}_1$. Consider a vector bundle E_1 on C_1 that is the direct sum of h indecomposable generic vector bundles of rank $\bar{n}$ and degree $\bar{d}_1$. Take a generic vector bundle E_g on C_g of rank n and degree d_g . Glue E_1, E_g at P by a generic gluing. The resulting vector bundle E is a generic point in one of the components of the moduli space of vector bundles of rank n and degree d on C (cf [T1] Theorem).

We first show that $A(n', d, E)$ is non-empty and has a component of dimension $s - n'(n-n')g$. Define $d'_g = d' - d'_1$. Then, $n'd_g - nd'_g = s - n'(n-n') \geq n'(n-n')(g-1)$. By (0,1), $A_g(n', d_g, E_g)$ is non-empty and has a component of $\dim n'd_g - nd'_g - n'(n-n')(g-1)$. The map to $\mathbf{G}(n', E_{g, P_g})$ is either onto or it has finite fibers. By (2,8), $A_1(n', d_1, E_1)$ is non-empty and has a component of $\dim n'd_1 - nd'_1 = n'(n-n')$. The map to $\mathbf{G}(n', E_{1, P_1})$ is onto and has finite fibers. Therefore, each of the subbundles of degree d'_g of E_g glues with a finite number of subbundles of degree d'_1 of E_1 . This shows that the component $A(n', (d'_1, d'_g), E)$ of $A(n', d, E)$ (corresponding to the given splitting of the degree of the subbundle between the two components) is non-empty of dimension $n'd_g - nd'_g - n'(n-n')(g-1) = s - n'(n-n')g$. From (0,1), (2,8) and (2,6), stability conditions are satisfied for the restrictions to the components. From [T1], they give rise to a stable E .

We need to check that E does not have higher dimensional varieties of subbundles of rank n' and degree d' corresponding to a different splitting of the degree $d' = \bar{d}'_1 + \bar{d}'_g$. Assume that $A(n', (\bar{d}'_1, \bar{d}'_g), E)$ is non-empty. By the genericity of E_g , $n'd_g - n\bar{d}'_g \geq n'(n-n')(g-1)$ (cf [L]). By (0,1), $\dim A(n', \bar{d}'_g, E_g) = n'd_g - n\bar{d}'_g - n'(n-n')(g-1)$ and either is mapped by π_{P_g} onto $\mathbf{G}(n', E_{g, P_g})$ or the map has finite fibers. By (2,8), $\dim A(n', \bar{d}'_1, E_1) = n'd_1 - n\bar{d}'_1 - n'(n-n')(g-1)$ and either is mapped by π_{P_1} onto $\mathbf{G}(n', E_{1, P_1})$ or the map has finite fibers. An element in $A(n', (\bar{d}'_1, \bar{d}'_g), E)$ is determined by an element in $A(n', \bar{d}'_1, E_1)$ and an element in $A(n', \bar{d}'_g, E_g)$ whose images in $\mathbf{G}(n', E_{1, P_1})$ and $\mathbf{G}(n', E_{g, P_g})$ glue together.

If π_{P_g} is onto, then each element in $A(n', \bar{d}'_1, E_1)$ glues with a fiber of π_{P_g} . Hence we obtain a set of dimension $n'd_1 - n\bar{d}'_1 + (n'd_g - n\bar{d}'_g - n'(n-n')) = s - n'(n-n')g$ as desired. If π_{P_1} is onto, one can reason in the same way. Assume now that both maps have finite fibers. Then by (3,1), the genericity of the gluing implies that $n'd_g - n\bar{d}'_g - n'(n-n') + n'd_1 - n\bar{d}'_1 + n^2 - n'(n-n') \geq n^2$. This can be written as $s \geq n'(n-n')g$. Then, by (3,1)iv) the genericity of the gluing and the finiteness of the fibers of the maps π_{P_1}, π_{P_g} ,

$$\dim A(n', (\bar{d}_1, \bar{d}_g), E) = \dim A_g(n', \bar{d}_g, E_g) + \dim A_1(n', \bar{d}_1, E_1) - n'(n-n')g = s - n'(n-n')g.$$

Consider now the fiber F of $\pi_{Q|A_1(n', \bar{d}_1, E_1)}$ over a generic point of $\mathbf{G}(n', E_{1, P_Q})$. From (2,8), either π_{P_1} maps F onto $\mathbf{G}(n', E_{1, P_1})$ or it maps with finite fibers. As $A_g(n', \bar{d}_g, E_g)$ either maps onto $\mathbf{G}(n', E_{P_g})$ or it maps with finite fibers, the result follows with the same line of reasoning as before.

Consider now the case $g = 1$. We can repeat the argument above replacing (0,1) by (2,8) except that we have semistability instead of stability for the restriction of the vector bundle to $C_g = C'_1$. From the remark at the end of (1,2), we need to prove that the resulting vector bundle E on the reducible curve is stable for some polarisation. If this were not the case, there would be a subbundle F of E with the same rank on each component and such that $\mu(F) = \mu(E)$ (cf [T1], end of the proof of step 2, p.436). This latter condition along with the semistability of each restriction implies that $\mu(F|_{C_1}) = \mu(E|_{C_1}), \mu(F|_{C_2}) = \mu(E|_{C_2})$. There are only a finite number of subbundles F'_1, F_1 satisfying these conditions. From (3,1), the dimension of the set of identifications of E_{1, P_1} with E'_{1, P'_1} that glue together a pair of these subspaces is at most $n^2 - n'(n-n')$. This is smaller than the dimension of the X considered above. This finishes the proof.

The proof of (0,1) can now be carried out by induction on g using (1,2) and (3,2).

References.

- [A] M.Atiyah, Vector bundles on elliptic curves, Proc.London Math.Soc. **(3)7**(1957),414-452.
- [B,B,R] E.Ballico, L.Brambila-Paz, B.Russo. Exact sequences of stable vector bundles on projective curves, Math.Nachr.
- [L]H.Lange, Zur Klassifikation von Regelmanigfaltigkeiten, Math.Ann. **262**(1983), 447-459.
- [L2] H.Lange, Universal Families of Extensions, J. of Algebra **83** (1983), 101-112.
- [L,N] H.Lange, M.S.Narasimhan, Maximal subbundles of rank two vector bundles on curves, Math.Ann. **266** (1983), 55-72.
- [P] R.Pandharipande, A compactification over $\bar{\mathcal{M}}_g$ of the universal moduli space of slope-semistable vector bundles, Journal of the A.M.S. **V9 N2**(1996), 425-471.
- [S] C.S.Seshadri. Fibres vectoriels sur les courbes algebriques. Asterisque **96**, Societe Mathematique de France, 1982.
- [T1]M. Teixidor i Bigas, Moduli spaces of semistable bundles on tree-like curves, Math.Ann.**290**(1991), 341-348.
- [T2]M. Teixidor i Bigas, Moduli spaces of semistable bundles on reducible curves, Amer.J.of Math. **117** (1995), 125-139
- [T3]M. Teixidor i Bigas, Stable extensions by line bundles, preprint Jan. 97.
- [Tu]L.Tu, Semistable bundles over an elliptic curve, Adv in Math **98**(1993),1-26.